

# American Board of Ophthalmology Office Record Review Module

## Diagnosis/Case: Stable Primary Open-Angle Glaucoma

**Module Definition:** The patient must be 18 years or older and the patient's care must have been initiated and followed by you for a minimum of two years, and seen by you at least once in the past 12 months. The patient's glaucoma must fit into the following status:

- Stable, mild to moderate disc and/or visual field damage (no more than nasal step or arcuate scotoma in one hemisphere).
- Therapeutic adjustments may have been made, but none within the past year.

History (Positive or Negative Findings)			
Recorded	Not Recorded	Not Applicable	Item
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	1. Reason for visit/chief complaint
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	2. Ocular history/ symptoms
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	3. Family history of glaucoma
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	4. Risk factors for glaucoma
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	5. Systemic disorders
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	6. Ocular medications
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	7. Systemic medications
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	8. Drug reactions/ allergies

Examination and Diagnostic Procedures			
Recorded	Not Recorded	Not Applicable	Item
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	1. Visual acuity
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	2. Pupillary examination
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	3. Intraocular pressures (IOP)
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	4. Corneal thickness
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	5. Slit-lamp biomicroscopy
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	6. Gonioscopy
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	7. Fundus examination
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	8. Optic nerve evaluation
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	9. Visual fields
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	10. Scanning laser evaluation of optic disc and RNFL

Diagnostic Techniques			
Recorded	Not	Not	Item

	Recorded	Applicable	
			Tonometer used to measure IOP:
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	1. Goldmann applanation
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	2. TonoPen
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	3. Other
			Gonioscopy lens used:
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	4. Single or 3-mirror (Goldmann)
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	5. 4-mirror (Zeiss/Posner/Sussman)
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	6. Koepe (direct gonioscopy)
			Technique used for disc/fundus examination:
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	7. Direct ophthalmoscope
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	8. Indirect slit-lamp lenses (90D lens, etc.)
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	9. Hruby lens
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	10. Other
			Method used to document visual fields:
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	11. Automated, bowl
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	12. Manual, bowl
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	13. Short-wavelength automated perimetry (SWAP)
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	14. Other

### Assessment and Management

Recorded	Not Recorded	Not Applicable	Item
<input type="radio"/>	<input type="radio"/>		1. Current diagnosis
<input type="radio"/>	<input type="radio"/>		2. Target pressure or range: reduction of IOP by minimum of 15%, or documentation of a plan of care if this target is not achieved
<input type="radio"/>	<input type="radio"/>		3. Initial assessment of glaucoma severity
<input type="radio"/>	<input type="radio"/>		4. Current treatment plan
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	5. Discussion of potential risks/side effects of treatment medications
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	6. Instructions provided for use of medication
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	7. Informed consent for treatment procedure
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	8. IOP measurement 30-120 minutes after laser trabeculoplasty
<input type="radio"/>	<input type="radio"/>		9. Counseling on glaucoma in the past year

### Management Techniques

Recorded	Not Recorded	Not Applicable	Item
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	1. Beta-adrenergic antagonists
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	2. Prostaglandins
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	3. Parasympathomimetics (cholinergic agonists, miotics)
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	4. Alpha- adrenergic agonists/epinephrine
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	5. Carbonic anhydrase inhibitors (topical)

<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	6. Carbonic anhydrase inhibitors (systemic) (oral or IV)
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	7. Argon laser trabeculoplasty
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	8. Selective laser trabeculoplasty
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	9. Filtering surgery
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	10. Cyclodestruction
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	11. Monocular trial of eye medications

Follow-Up			
Recorded	Not Recorded	Not Applicable	Item
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	1. Reason for visit/interval history
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	2. Current ocular medication and therapeutic compliance
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	3. Side effects from medications
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	4. Pulse check (if on Beta-blockers)
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	5. Visual acuity
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	6. Intraocular pressures (IOP)
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	7. Slit-lamp examination
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	8. Optic disc appearance
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	9. Visual fields
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	10. Scanning laser evaluation of optic disc and RNFL

Follow-Up Intervals			
Recorded	Not Recorded	Not Applicable	Item
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	1. Follow-up after initiating/adjusting therapy
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	2. Follow-up after glaucoma stabilized
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	3. Visual fields examination
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	4. Gonioscopy
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	5. Optic evaluation
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	6. Pupillary dilation in the past 12 months

Safety and Environment			
Recorded	Not Recorded	Not Applicable	Item
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	1. Tonometer tip covered and cleaned after each patient contact
<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	2. Periodic tonometer and Snellen chart calibration